

GRAPHS WITHOUT REPEATED CYCLE LENGTHS

Chunhui Lai *

Department of Mathematics, Zhangzhou Teachers College, Zhangzhou,
Fujian 363000, P. R. of CHINA; and Graph Theory and Combinatorics Laboratory,
Institute of Systems Science, Academy of Mathematics and Systems Science,
Chinese Academy of Sciences, Beijing 100080, P. R. of CHINA

`zjlaichu@public.zzptt.fj.cn`

MR Subject Classifications: 05C38, 05C35

Key words: graph, cycle, number of edges

Abstract

In 1975, P. Erdős proposed the problem of determining the maximum number $f(n)$ of edges in a graph of n vertices in which any two cycles are of different lengths. In this paper, it is proved that

$$f(n) \geq n + 36t$$

for $t = 1260r + 169$ ($r \geq 1$) and $n \geq 540t^2 + \frac{175811}{2}t + \frac{7989}{2}$. Consequently,
 $\liminf_{n \rightarrow \infty} \frac{f(n)-n}{\sqrt{n}} \geq \sqrt{2 + \frac{2}{5}}$.

1 Introduction

Let $f(n)$ be the maximum number of edges in a graph on n vertices in which no two cycles have the same length. In 1975, Erdős raised the problem of determining $f(n)$ (see [1], p.247, Problem 11). Shi[2] proved that

$$f(n) \geq n + [(\sqrt{8n-23} + 1)/2]$$

for $n \geq 3$. Lai[3,4,5,6,7] proved that for $n \geq \frac{6911}{16}t^2 + \frac{514441}{8}t - \frac{3309665}{16}$, $t = 27720r + 169$,

$$f(n) \geq n + 32t - 1,$$

*Project Supported by NSF of Fujian(A96026), Science and Technology Project of Fujian(K20105) and Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering".

and for $n \geq e^{2m}(2m+3)/4$,

$$f(n) < n - 2 + \sqrt{n \ln(4n/(2m+3))} + 2n + \log_2(n+6).$$

Boros, Caro, Füredi and Yuster[8] proved that

$$f(n) \leq n + 1.98\sqrt{n}(1 + o(1)).$$

In this paper, we construct a graph G having no two cycles with the same length which leads the following result.

Theorem. Let $t = 1260r + 169$ ($r \geq 1$), then

$$f(n) \geq n + 36t$$

for $n \geq 540t^2 + \frac{175811}{2}t + \frac{7989}{2}$.

2 Proof of the theorem

Proof. Let $t = 1260r + 169, r \geq 1, n_t = 540t^2 + \frac{175811}{2}t + \frac{7989}{2}, n \geq n_t$. We shall show that there exists a graph G on n vertices with $n + 36t$ edges such that all cycles in G have distinct lengths.

Now we construct the graph G which consists of a number of subgraphs: B_i , $(0 \leq i \leq 21t - 1, 27t \leq i \leq 28t + 64, 29t - 734 \leq i \leq 29t + 267, 30t - 531 \leq i \leq 30t + 57, 31t - 741 \leq i \leq 31t + 58, 32t - 740 \leq i \leq 32t + 57, 33t - 741 \leq i \leq 33t + 57, 34t - 741 \leq i \leq 34t + 52, 35t - 746 \leq i \leq 35t + 60, 36t - 738 \leq i \leq 36t + 60, 37t - 738 \leq i \leq 37t + 799, i = 21t + 2j + 1 (0 \leq j \leq t - 1), i = 21t + 2j (0 \leq j \leq \frac{t-1}{2}), i = 23t + 2j + 1 (0 \leq j \leq \frac{t-1}{2}), \text{ and } i = 26t)$.

Now we define these B_i 's. These subgraphs all have a common vertex x , otherwise their vertex sets are pairwise disjoint.

For $0 \leq i \leq t - 1$, let the subgraph $B_{21t+2i+1}$ consist of a cycle

$$xu_i^1 u_i^2 \dots u_i^{25t+2i-1} x$$

and a path:

$$xu_{i,1}^1 u_{i,1}^2 \dots u_{i,1}^{(19t+2i-1)/2} u_i^{(23t+2i+1)/2}$$

Based the construction, $B_{21t+2i+1}$ contains exactly three cycles of lengths:

$$21t + 2i + 1, 23t + 2i, 25t + 2i.$$

For $0 \leq i \leq \frac{t-3}{2}$, let the subgraph B_{21t+2i} consist of a cycle

$$xv_i^1 v_i^2 \dots v_i^{25t+2i} x$$

and a path:

$$xv_{i,1}^1 v_{i,1}^2 \dots v_{i,1}^{9t+i-1} v_i^{12t+i}$$

Based the construction, B_{21t+2i} contains exactly three cycles of lengths:

$$21t + 2i, 22t + 2i + 1, 25t + 2i + 1.$$

For $0 \leq i \leq \frac{t-3}{2}$, let the subgraph $B_{23t+2i+1}$ consist of a cycle

$$xw_i^1 w_i^2 \dots w_i^{26t+2i+1} x$$

and a path:

$$xw_{i,1}^1 w_{i,1}^2 \dots w_{i,1}^{(21t+2i-1)/2} w_i^{(25t+2i+1)/2}$$

Based the construction, $B_{23t+2i+1}$ contains exactly three cycles of lengths:

$$23t + 2i + 1, 24t + 2i + 2, 26t + 2i + 2.$$

For $58 \leq i \leq t - 742$, let the subgraph $B_{27t+i-57}$ consist of a cycle

$$C_{27t+i-57} = xy_i^1 y_i^2 \dots y_i^{132t+11i+893} x$$

and ten paths sharing a common vertex x , the other end vertices are on the cycle $C_{27t+i-57}$:

$$\begin{aligned} & xy_{i,1}^1 y_{i,1}^2 \dots y_{i,1}^{(17t-1)/2} y_i^{(37t-115)/2+i} \\ & xy_{i,2}^1 y_{i,2}^2 \dots y_{i,2}^{(19t-1)/2} y_i^{(57t-103)/2+2i} \\ & xy_{i,3}^1 y_{i,3}^2 \dots y_{i,3}^{(19t-1)/2} y_i^{(77t+315)/2+3i} \\ & xy_{i,4}^1 y_{i,4}^2 \dots y_{i,4}^{(21t-1)/2} y_i^{(97t+313)/2+4i} \\ & xy_{i,5}^1 y_{i,5}^2 \dots y_{i,5}^{(21t-1)/2} y_i^{(117t+313)/2+5i} \\ & xy_{i,6}^1 y_{i,6}^2 \dots y_{i,6}^{(23t-1)/2} y_i^{(137t+311)/2+6i} \\ & xy_{i,7}^1 y_{i,7}^2 \dots y_{i,7}^{(23t-1)/2} y_i^{(157t+309)/2+7i} \\ & xy_{i,8}^1 y_{i,8}^2 \dots y_{i,8}^{(25t-1)/2} y_i^{(177t+297)/2+8i} \\ & xy_{i,9}^1 y_{i,9}^2 \dots y_{i,9}^{(25t-1)/2} y_i^{(197t+301)/2+9i} \\ & xy_{i,10}^1 y_{i,10}^2 \dots y_{i,10}^{(27t-1)/2} y_i^{(217t+305)/2+10i}. \end{aligned}$$

As a cycle with d chords contains $\binom{d+2}{2}$ distinct cycles, $B_{27t+i-57}$ contains exactly 66 cycles of lengths:

$27t + i - 57,$	$28t + i + 7,$	$29t + i + 210,$	$30t + i,$
$31t + i + 1,$	$32t + i,$	$33t + i,$	$34t + i - 5,$
$35t + i + 3,$	$36t + i + 3,$	$37t + i + 742,$	$38t + 2i - 51,$
$38t + 2i + 216,$	$40t + 2i + 209,$	$40t + 2i,$	$42t + 2i,$
$42t + 2i - 1,$	$44t + 2i - 6,$	$44t + 2i - 3,$	$46t + 2i + 5,$
$46t + 2i + 744,$	$48t + 3i + 158,$	$49t + 3i + 215,$	$50t + 3i + 209,$
$51t + 3i - 1,$	$52t + 3i - 1,$	$53t + 3i - 7,$	$54t + 3i - 4,$
$55t + 3i - 1,$	$56t + 3i + 746,$	$59t + 4i + 157,$	$59t + 4i + 215,$
$61t + 4i + 208,$	$61t + 4i - 2,$	$63t + 4i - 7,$	$63t + 4i - 5,$
$65t + 4i - 2,$	$65t + 4i + 740,$	$69t + 5i + 157,$	$70t + 5i + 214,$
$71t + 5i + 207,$	$72t + 5i - 8,$	$73t + 5i - 5,$	$74t + 5i - 3,$
$75t + 5i + 739,$	$80t + 6i + 156,$	$80t + 6i + 213,$	$82t + 6i + 201,$
$82t + 6i - 6,$	$84t + 6i - 3,$	$84t + 6i + 738,$	$90t + 7i + 155,$
$91t + 7i + 207,$	$92t + 7i + 203,$	$93t + 7i - 4,$	$94t + 7i + 738,$
$101t + 8i + 149,$	$101t + 8i + 209,$	$103t + 8i + 205,$	$103t + 8i + 737,$
$111t + 9i + 151,$	$112t + 9i + 211,$	$113t + 9i + 946,$	$122t + 10i + 153,$
$122t + 10i + 952,$	$132t + 11i + 894.$		

B_0 is a path with an end vertex x and length $n - n_t$. Other B_i is simply a cycle of length i .

Then $f(n) \geq n + 36t$, for $n \geq n_t$. This completes the proof.

From the above theorem, we have

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2 + \frac{2}{5}},$$

which is better than the previous bounds $\sqrt{2}$ (see [2]), $\sqrt{2 + \frac{2562}{6911}}$ (see [7]).

Combining this with Boros, Caro, Füredi and Yuster's upper bound, we get

$$1.98 \geq \limsup_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2.4}.$$

We make the following conjecture:

Conjecture.

$$\lim_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} = \sqrt{2.4}.$$

Acknowledgment

The author thanks Prof. Yair Caro and Raphael Yuster for sending me references[8]. The author thanks Prof. Genghua Fan and Cheng Zhao for their valuable suggestions.

References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Macmillan, New York, 1976).
- [2] Y. Shi, On maximum cycle-distributed graphs, Discrete Math. 71(1988) 57-71.
- [3] Chunhui Lai, On the Erdős problem, J. Zhangzhou Teachers College(Natural Science Edition) 3(1)(1989) 55-59.
- [4] Chunhui Lai, Upper bound and lower bound of $f(n)$, J. Zhangzhou Teachers College(Natural Science Edition) 4(1)(1990) 29,30-34.
- [5] Chunhui Lai, On the size of graphs with all cycle having distinct length, Discrete Math. 122(1993) 363-364.
- [6] Chunhui Lai, The number of edges in a graph in which no two cycles have the same length, J. Zhangzhou Teachers College (Natural Science Edlition) 8(4)(1994), 30-34.
- [7] Chunhui Lai, A lower bound for the number of edges in a graph containing no two cycles of the same length, The Electronic J. of Combinatorics 8(2001), #N9.
- [8] E. Boros, Y. Caro, Z. Füredi and R. Yuster, Covering non-uniform hypergraphs (submitted, 2000).